

## GASTRIC CANCER TREATMENT REGIMENS (Part 1 of 7)

**Clinical Trials:** The NCCN recommends cancer patient participation in clinical trials as the gold standard for treatment.

Cancer therapy selection, dosing, administration, and the management of related adverse events can be a complex process that should be handled by an experienced healthcare team. Clinicians must choose and verify treatment options based on the individual patient; drug dose modifications and supportive care interventions should be administered accordingly. The cancer treatment regimens below may include both U.S. Food and Drug Administration-approved and unapproved indications/regimens. These regimens are only provided to supplement the latest treatment strategies.

These Guidelines are a work in progress that may be refined as often as new significant data becomes available. The NCCN Guidelines® are a consensus statement of its authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult any NCCN Guidelines® is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network makes no warranties of any kind whatsoever regarding their content, use, or application and disclaims any responsibility for their application or use in any way.

### Principles of Systemic Therapy<sup>1</sup>

Chemotherapy regimens should be chosen in the context of performance status, medical comorbidities, toxicity profile, and HER2-neu expression (for adenocarcinoma only). Two-drug cytotoxic regimens are preferred for patients with advanced disease because of lower toxicity. Three-drug cytotoxic regimens should be reserved for medically fit patients with good performance status and access to frequent toxicity evaluation. Doses and schedules for any regimen that is not derived from category 1 evidence is a suggestion, and subject to appropriate modifications depending on the circumstances. Infusional fluorouracil and capecitabine may be used interchangeably (except as indicated). Cisplatin and oxaliplatin may be used interchangeably depending on toxicity profile.

### Preoperative Chemoradiation (esophagogastric junction and gastric cardia)<sup>1</sup>

**Note:** All recommendations are Category 2A unless otherwise indicated.

#### Preferred Regimens

REGIMEN	DOSING
<b>Paclitaxel + carboplatin (Category 1)<sup>2</sup></b>	<b>Day 1:</b> Paclitaxel 50mg/m <sup>2</sup> IV + carboplatin AUC 2mg/mL × min IV. Repeat weekly for 5 weeks.
<b>Cisplatin + 5-fluorouracil (5-FU) (Category 1)<sup>3,4</sup></b>	<b>Days 1 and 29:</b> Cisplatin 75–100mg/m <sup>2</sup> IV <b>Days 1–4 and 29–32:</b> 5-FU 750–1000mg/m <sup>2</sup> IV continuous infusion over 24 hours. <b>OR</b> <b>Days 1–5:</b> Cisplatin 15mg/m <sup>2</sup> IV once daily plus 5-FU 800mg/m <sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 21 days for 2 cycles.
<b>Oxaliplatin + 5-FU (Category 1)<sup>5,6</sup></b>	<b>Days 1, 15, and 29:</b> Oxaliplatin 85mg/m <sup>2</sup> IV <b>Days 1–33:</b> 5-FU 180mg/m <sup>2</sup> IV. <b>OR</b> <b>Day 1:</b> Oxaliplatin 85mg/m <sup>2</sup> and leucovorin 400mg/m <sup>2</sup> <b>followed by</b> 5-FU 400mg/m <sup>2</sup> bolus, then 800mg/m <sup>2</sup> 24-hour continuous infusion over days 1 and 2; the first 3 cycles were delivered during radiotherapy (RT), the other 3 after RT; 6 bimonthly (14 days) cycles.
<b>Cisplatin + capecitabine<sup>7</sup></b>	<b>Day 1:</b> Cisplatin 30mg/m <sup>2</sup> IV <b>Days 1–5:</b> Capecitabine 800mg/m <sup>2</sup> orally twice daily. Repeat cycle weekly for 5 weeks.
<b>Oxaliplatin + capecitabine<sup>8</sup></b>	<b>Days 1, 15, and 29:</b> Oxaliplatin 85mg/m <sup>2</sup> IV <b>Days 1–5:</b> Capecitabine 625mg/m <sup>2</sup> orally twice daily for 5 weeks.

#### Other Regimens

<b>Irinotecan + cisplatin (Category 2B)<sup>9</sup></b>	<b>Days 1, 8, 22, 29:</b> Irinotecan 65mg/m <sup>2</sup> IV plus cisplatin 30mg/m <sup>2</sup> IV.
<b>Paclitaxel + 5-FU (Category 2B)<sup>10</sup></b>	<b>Day 1:</b> Paclitaxel 45–50mg/m <sup>2</sup> IV weekly <b>Days 1–5:</b> 5-FU 300mg/m <sup>2</sup> IV continuous infusion. Weekly for 5 weeks.
<b>Paclitaxel + capecitabine (Category 2B)<sup>10</sup></b>	<b>Day 1:</b> Paclitaxel 45–50mg/m <sup>2</sup> IV <b>Days 1–5:</b> Capecitabine 625–825mg/m <sup>2</sup> orally BID. Weekly for 5 weeks.

### Perioperative Chemotherapy (including esophagogastric junction)<sup>1</sup>

<b>Epirubicin + cisplatin + 5-FU (ECF) (Category 1)<sup>11</sup></b>	<b>Day 1:</b> Epirubicin 50mg/m <sup>2</sup> IV bolus + cisplatin 60mg/m <sup>2</sup> IV <b>Days 1–21:</b> 5-FU 200mg/m <sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 21 days for 3 cycles preoperatively and 3 cycles postoperatively.
--	--

*continued*

## GASTRIC CANCER TREATMENT REGIMENS (Part 2 of 7)

### Perioperative Chemotherapy (including esophagogastric junction)<sup>1</sup> (continued)

REGIMEN	DOSING
<b>ECF modification: epirubicin + oxaliplatin + 5-FU<sup>12</sup></b>	<b>Day 1:</b> Epirubicin 50mg/m <sup>2</sup> IV; oxaliplatin 130mg/m <sup>2</sup> IV <b>Days 1–21:</b> 5-FU 200mg/m <sup>2</sup> IV continuous infusion over 24 hours. Repeat cycle every 21 days for 3 cycles preoperatively and 3 cycles postoperatively.
<b>ECF modification: epirubicin + cisplatin + capecitabine<sup>12</sup></b>	<b>Day 1:</b> Epirubicin 50mg/m <sup>2</sup> IV; cisplatin 60mg/m <sup>2</sup> IV <b>Days 1–21:</b> Capecitabine 625mg/m <sup>2</sup> orally twice daily. Repeat cycle every 21 days for 3 cycles preoperatively and 3 cycles postoperatively.
<b>ECF modification: epirubicin + oxaliplatin + capecitabine<sup>12</sup></b>	<b>Day 1:</b> Epirubicin 50mg/m <sup>2</sup> IV; oxaliplatin 130mg/m <sup>2</sup> IV <b>Days 1–21:</b> Capecitabine 625mg/m <sup>2</sup> orally twice daily. Repeat cycle every 21 days for 3 cycles preoperatively and 3 cycles postoperatively.
<b>5-FU + cisplatin (Category 1)<sup>13</sup></b>	<b>Day 1:</b> Cisplatin 75–80mg/m <sup>2</sup> IV <b>Days 1–5:</b> 5-FU 800mg/m <sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 28 days for 2–3 cycles preoperatively and 3–4 cycles postoperatively for a total of 6 cycles.

### Postoperative Chemoradiation (including esophagogastric junction)<sup>1</sup>

<b>5-FU + leucovorin<sup>14</sup></b>	<b>Cycles 1, 3, and 4 (before and after radiation)</b> <b>Days 1–5:</b> Leucovorin 20mg/m <sup>2</sup> IVP plus 5-FU 425mg/m <sup>2</sup> /day IVP Repeat cycle every 28 days. <b>Cycle 2 (with radiation)</b> <b>Days 1–4 and 31–33:</b> Leucovorin 20mg/m <sup>2</sup> IVP <b>Days 1–4:</b> 5-FU 400mg/m <sup>2</sup> /day IVP. Repeat cycle every 35 days.  The NCCN panel acknowledges that the Intergroup 0116 Trial formed the basis for post-operative adjuvant chemoradiation strategy. However, the panel does not recommend the above specified doses or schedule of cytotoxic agents because of concerns regarding toxicity. The panel recommends one of the following modifications instead.
<b>Capecitabine<sup>15</sup></b>	<b>Days 1–14:</b> Capecitabine 750–1000mg/m <sup>2</sup> orally twice daily. Repeat cycle every 28 days; 1 cycle before and 2 cycles after chemoradiation.
<b>5-FU + leucovorin<sup>16</sup></b>	<b>Days 1, 2, 15, and 16:</b> Leucovorin 400mg/m <sup>2</sup> IV followed by 5-FU 400mg/m <sup>2</sup> IVP and a 24-hour infusion of 5-FU 1200mg/m <sup>2</sup> ; 1 cycle before and 2 cycles after chemoradiation. Repeat cycle every 28 days.
<b>5-FU with radiation<sup>17</sup></b>	<b>Days 1–5 OR Days 1–7:</b> 5-FU 200–250mg/m <sup>2</sup> IV continuous infusion over 24 hours once daily; weekly for 5 weeks.
<b>Capecitabine with radiation<sup>18</sup></b>	<b>Days 1–5 OR Days 1–7:</b> Capecitabine 625–825mg/m <sup>2</sup> orally twice daily; weekly for 5 weeks.

### Postoperative Chemotherapy

<b>Capecitabine + oxaliplatin<sup>19</sup></b>	<b>Days 1–14:</b> Capecitabine 1000mg/m <sup>2</sup> orally twice daily <b>Day 1:</b> Oxaliplatin 130mg/m <sup>2</sup> IV. Repeat cycle every 21 days for 8 cycles.
<b>Capecitabine + cisplatin<sup>20</sup></b>	<b>Days 1–14:</b> Capecitabine 1000mg/m <sup>2</sup> orally twice daily <b>Day 1:</b> Cisplatin 60mg/m <sup>2</sup> IV. Repeat cycle every 21 days for 6 cycles.

### Metastatic or Locally Advanced Cancer (where local therapy is not indicated)<sup>1</sup>

#### First-line Therapy

<b>Trastuzumab + chemotherapy (NOTE: for HER2-neu overexpressing adenocarcinoma)<sup>21</sup></b>	<b>Day 1:</b> Trastuzumab 8mg/kg IV loading dose (Cycle 1 only); <b>followed by</b> trastuzumab 6mg/kg IV every 3 weeks, plus chemotherapy <b>OR</b> <b>Day 1 of Cycle 1:</b> Trastuzumab 6mg/kg IV loading dose, then 4mg/kg IV every 14 days. <b>Chemotherapy:</b> <b>Day 1:</b> Cisplatin 80mg/m <sup>2</sup> IV, <b>plus</b> <b>Days 1–14:</b> Capecitabine 1000mg/m <sup>2</sup> orally twice daily. (Category 1) <b>OR</b> <b>Days 1–5:</b> 5-FU 800mg/m <sup>2</sup> continuous IV infusion. Repeat cycle every 21 days for 6 cycles
---	---

*continued*

## GASTRIC CANCER TREATMENT REGIMENS (Part 3 of 7)

### Metastatic or Locally Advanced Cancer (where local therapy is not indicated)<sup>1</sup> (continued)

#### First-line Therapy (continued)

REGIMEN	DOSING
<b>Docetaxel + cisplatin + 5-FU (DCF) (Category 1)</b> <sup>22</sup>	<p><b>Day 1:</b> Docetaxel 75mg/m<sup>2</sup> IV + cisplatin 75mg/m<sup>2</sup> IV</p> <p><b>Days 1-5:</b> 5-FU 1,000mg/m<sup>2</sup> IV continuous infusion over 24 hours, daily. Repeat cycle every 28 days.</p> <p>The panel does not recommend the above specified doses or schedule of cytotoxic agents because of concerns regarding toxicity.</p>
<b>Modified DCF (docetaxel + cisplatin + leucovorin + 5-FU)</b> <sup>23</sup>	<p><b>Day 1:</b> Docetaxel 40mg/m<sup>2</sup> IV plus leucovorin 400mg/m<sup>2</sup> IV plus 5-FU 400mg/m<sup>2</sup></p> <p><b>Days 1 and 2:</b> 5-FU 1000mg/m<sup>2</sup> IV continuous infusion over 24 hours</p> <p><b>Day 3:</b> Cisplatin 40mg/m<sup>2</sup> IV.</p> <p>Repeat cycle every 14 days.</p>
<b>Modified DCF (docetaxel + oxaliplatin + 5-FU)</b> <sup>24</sup>	<p><b>Day 1:</b> Docetaxel 50mg/m<sup>2</sup> IV plus oxaliplatin 85mg/m<sup>2</sup></p> <p><b>Days 1 and 2:</b> 5-FU 1,200mg/m<sup>2</sup> IV continuous infusion over 24 hours. Repeat cycle every 14 days.</p>
<b>Modified DCF (docetaxel + carboplatin + 5-FU) (Category 2B)</b> <sup>25</sup>	<p><b>Day 1:</b> Docetaxel 75mg/m<sup>2</sup></p> <p><b>Day 2:</b> Carboplatin AUC 6mg/mL × min</p> <p><b>Days 1-3:</b> 5-FU 1,200mg/m<sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 21 days.</p>
<b>ECF (Category 1)</b> <sup>26</sup>	<p><b>Day 1:</b> Epirubicin 50mg/m<sup>2</sup> IV bolus + cisplatin 60mg/m<sup>2</sup> IV</p> <p><b>Days 1-21:</b> 5-FU 200mg/m<sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 21 days.</p>
<b>ECF modifications (epirubicin + oxaliplatin + 5-FU) (Category 1)</b> <sup>27</sup>	<p><b>Days 1:</b> Epirubicin 50mg/m<sup>2</sup> IV plus oxaliplatin 130mg/m<sup>2</sup> IV</p> <p><b>Days 1-21:</b> 5-FU 200mg/m<sup>2</sup> IV continuous infusion over 24 hours. Repeat cycle every 21 days.</p>
<b>ECF modifications (epirubicin + cisplatin + capecitabine) (Category 1)</b> <sup>27</sup>	<p><b>Day 1:</b> Epirubicin 50mg/m<sup>2</sup> IV plus cisplatin 60mg/m<sup>2</sup> IV</p> <p><b>Days 1-21:</b> Capecitabine 625mg/m<sup>2</sup> orally twice daily. Repeat cycle every 21 days.</p>
<b>ECF modifications (epirubicin + oxaliplatin + capecitabine) (Category 1)</b> <sup>27</sup>	<p><b>Day 1:</b> Epirubicin 50mg/m<sup>2</sup> IV plus oxaliplatin 130mg/m<sup>2</sup> IV</p> <p><b>Days 1-21:</b> capecitabine 625mg/m<sup>2</sup> IV orally twice daily. Repeat cycle every 21 days.</p>
<b>Fluoropyrimidine and cisplatin (5-FU + cisplatin) (Category 1)</b> <sup>33</sup>	<p><b>Day 1:</b> Cisplatin 75-100mg/m<sup>2</sup> IV</p> <p><b>Days 1-4:</b> 5-FU 1,000mg/m<sup>2</sup> IV continuous infusion over 24 hours daily.</p>
<b>Fluoropyrimidine and cisplatin (5-FU + cisplatin + leucovorin) (Category 1)</b> <sup>28-31</sup>	<p><b>Day 1:</b> Cisplatin 50mg/m<sup>2</sup> IV plus leucovorin 200mg/m<sup>2</sup> IV plus 5-FU 2,000mg/m<sup>2</sup> IV continuous infusion over 24 hours. Repeat cycle every 14 days.</p>
<b>Fluoropyrimidine and cisplatin (capecitabine + cisplatin) (Category 1)</b> <sup>28-31</sup>	<p><b>Day 1:</b> Cisplatin 80mg/m<sup>2</sup> IV</p> <p><b>Day 1-14:</b> Capecitabine 1000mg/m<sup>2</sup> orally twice daily. Repeat cycle every 3 weeks.</p>
<b>Fluoropyrimidine and oxaliplatin (oxaliplatin + capecitabine)</b> <sup>29,32,33</sup>	<p><b>Day 1:</b> Oxaliplatin 130mg/m<sup>2</sup> IV</p> <p><b>Days 1-14:</b> Capecitabine 1000mg/m<sup>2</sup> orally twice daily. Repeat cycle every 21 days.</p>
<b>Fluoropyrimidine and oxaliplatin (oxaliplatin + leucovorin + 5-FU)</b> <sup>29,32,33</sup>	<p><b>Day 1:</b> Oxaliplatin 85mg/m<sup>2</sup> IV plus leucovorin 400mg/m<sup>2</sup> IV plus 5-FU 400mg/m<sup>2</sup> IVP</p> <p><b>Days 1 and 2:</b> 5-FU 1200mg/m<sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 14 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Day 1:</b> Oxaliplatin 85mg/m<sup>2</sup> IV plus leucovorin 200mg/m<sup>2</sup> IV plus 5-FU 2,600mg/m<sup>2</sup> IV continuous infusion over 24 hours. Repeat cycle every 14 days.</p>

*continued*

## GASTRIC CANCER TREATMENT REGIMENS (Part 4 of 7)

### Metastatic or Locally Advanced Cancer (where local therapy is not indicated)<sup>1</sup> (continued)

#### Other Regimens

REGIMEN	DOSING
<b>Fluorouracil and irinotecan (irinotecan + leucovorin + 5-FU) (Category 1)</b> <sup>34-36</sup>	<p><b>Day 1:</b> Irinotecan 80mg/m<sup>2</sup> IV plus leucovorin 500mg/m<sup>2</sup> IV plus 5-FU 2000mg/m<sup>2</sup> IV continuous infusion over 24 hours. Weekly for 6 weeks followed by 1 week off treatment <b>OR</b> weekly for 6 weeks followed by 2 weeks off treatment.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Day 1:</b> Irinotecan 150mg/m<sup>2</sup> IV plus leucovorin 20mg/m<sup>2</sup> IV <b>Days 1-2:</b> 5-FU 1000mg/m<sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 14 days.</p>
<b>Paclitaxel + cisplatin or carboplatin</b> <sup>37-39</sup>	<p><b>Day 1:</b> Paclitaxel 135-200mg/m<sup>2</sup> IV <b>Day 2:</b> Cisplatin 75mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Day 1:</b> Paclitaxel 90mg/m<sup>2</sup> IV plus cisplatin 50mg/m<sup>2</sup> IV. Repeat cycle every 14 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Day 1:</b> Paclitaxel 200mg/m<sup>2</sup> IV plus carboplatin AUC 5mg/mL × min Repeat cycle every 21 days.</p>
<b>Docetaxel + cisplatin</b> <sup>40,41</sup>	<p><b>Day 1:</b> Docetaxel 70-85mg/m<sup>2</sup> IV plus cisplatin 70-75mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p>
<b>Docetaxel + irinotecan (Category 2B)</b> <sup>42</sup>	<p><b>Days 1 and 8:</b> Docetaxel 35mg/m<sup>2</sup> IV plus irinotecan 50mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p>
<b>Fluoropyridimine</b> <sup>32,43,44</sup>	<p><b>Day 1:</b> Leucovorin 400mg/m<sup>2</sup> IV plus 5-FU 400mg/m<sup>2</sup> IVP <b>Days 1 and 2:</b> 5-FU 1200mg/m<sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 14 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Days 1-5:</b> 5-FU 800mg/m<sup>2</sup> IV continuous infusion over 24 hours daily. Repeat cycle every 28 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Days 1-14:</b> Capecitabine 1000-1250mg/m<sup>2</sup> orally twice daily. Repeat cycle every 21 days.</p>
<b>Taxane</b> <sup>45-48</sup>	<p><b>Day 1:</b> Docetaxel 75-100mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Day 1:</b> Paclitaxel 135-250mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Days 1, 8, 15 and 22:</b> Paclitaxel 80mg/m<sup>2</sup> IV once weekly. Repeat cycle every 28 days.</p>
<b>Second-line Therapy (preferred regimens)</b>	
<b>Ramucirumab (Category 1)</b> <sup>49</sup>	<p><b>Day 1:</b> Ramucirumab 8mg/kg IV. Repeat cycle every 14 days.</p>
<b>Ramucirumab + paclitaxel (Category 1)</b> <sup>50</sup>	<p><b>Day 1 and 15:</b> Ramucirumab 8mg/kg IV <b>Day 1, 8, and 15:</b> Paclitaxel 80mg/m<sup>2</sup>. Repeat cycle every 28 days.</p>
<b>Docetaxel (Category 1)</b> <sup>49</sup>	<p><b>Day 1:</b> Docetaxel 75-100mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p>
<b>Paclitaxel (Category 1)</b> <sup>50-52</sup>	<p><b>Day 1:</b> Paclitaxel 135-250mg/m<sup>2</sup> IV. Repeat cycle every 21 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Day 1:</b> Paclitaxel 80mg/m<sup>2</sup> IV once weekly. Repeat cycle every 28 days.</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Days 1, 8, and 15:</b> Paclitaxel 80mg/m<sup>2</sup> IV. Repeat cycle every 28 days.</p>

*continued*

## GASTRIC CANCER TREATMENT REGIMENS (Part 5 of 7)

### Metastatic or Locally Advanced Cancer (where local therapy is not indicated)<sup>1</sup> (continued)

#### Second-line Therapy (preferred regimens) (continued)

REGIMEN	DOSING
<b>Irinotecan (Category 1)</b> <sup>35,51-53</sup>	<b>Day 1:</b> Irinotecan 250–350mg/m <sup>2</sup> IV. Repeat cycle every 21 days. <b>OR</b> <b>Day 1:</b> Irinotecan 150–180mg/m <sup>2</sup> IV. Repeat cycle every 14 days. <b>OR</b> <b>Days 1 and 8:</b> Irinotecan 125mg/m <sup>2</sup> IV. Repeat cycle every 21 days.
<b>Other Regimens</b>	
<b>Irinotecan + cisplatin</b> <sup>32,54</sup>	<b>Days 1 and 8:</b> Irinotecan 65mg/m <sup>2</sup> IV plus cisplatin 25–30mg/m <sup>2</sup> IV. Repeat cycle every 21 days.
<b>Irinotecan + fluoropyridimine (Category 2B)</b> <sup>56,57</sup>	<b>Day 1:</b> Irinotecan 250mg/m <sup>2</sup> IV <b>Days 1–14:</b> Capecitabine 1000mg/m <sup>2</sup> PO BID daily. Repeat cycle every 21 days.
<b>Irinotecan + fluoropyridimine (Category 2B)</b> <sup>56,57</sup>	<b>Day 1:</b> Irinotecan 180mg/m <sup>2</sup> IV plus leucovorin 400mg/m <sup>2</sup> IV plus 5-FU 400mg/m <sup>2</sup> IVP <b>Days 1 and 2:</b> 5-FU 600–1200mg/m <sup>2</sup> /day IV continuous infusion on Days 1 and 2. Repeat cycle every 14 days.
<b>Docetaxel + irinotecan (Category 2B)</b> <sup>42</sup>	<b>Days 1 and 8:</b> Docetaxel 35mg/m <sup>2</sup> IV plus irinotecan 50mg/m <sup>2</sup> IV. Repeat cycle every 21 days.

#### Alternative Regimens for Consideration

<b>Mitomycin + irinotecan</b> <sup>57-59</sup>	<b>Day 1:</b> Mitomycin 6mg/m <sup>2</sup> IV <b>Days 2 and 9:</b> Irinotecan 125mg/m <sup>2</sup> Repeat cycle every 28 days. <b>OR</b> <b>Days 1 and 15:</b> Irinotecan 150mg/m <sup>2</sup> IV <b>Day 1:</b> Mitomycin 8mg/m <sup>2</sup> IV. Repeat cycle every 28 days. <b>OR</b> <b>Day 1:</b> Irinotecan 125mg/m <sup>2</sup> IV plus mitomycin 5mg/m <sup>2</sup> IV. Repeat cycle every 14 days.
<b>Mitomycin + leucovorin + 5-FU</b> <sup>60</sup>	<b>Days 1 and 22:</b> Mitomycin 10mg/m <sup>2</sup> IV <b>Day 1:</b> Leucovorin 500mg/m <sup>2</sup> IV plus 5-FU 2600mg/m <sup>2</sup> /day IV continuous infusion. Weekly for 6 weeks, followed by 2 weeks off treatment.

#### References

1. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology™. Gastric Cancer. v 3.2015. Available at: [http://www.nccn.org/professionals/physician\\_gls/pdf/gastric.pdf](http://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf). Accessed May 6, 2015.
2. van Hagen P, Hulshof MC, van Lanschot JJ, et al. CROSS Group. Preoperative chemoradiotherapy for esophageal or junctional cancer. *N Engl J Med*. 2012;366:2074–2084.
3. Tepper J, Krasna MJ, Niedzwiecki D, et al. Phase III trial of trimodality therapy with cisplatin, fluorouracil, radiotherapy, and surgery compared with surgery alone for esophageal cancer: CALGB 9781. *J Clin Oncol*. 2008;26:1086–1092.
4. Bedenne L, Michel P, Bouché O, et al. Chemoradiation followed by surgery compared with chemoradiation alone in squamous cancer of the esophagus: FFCD 9102. *J Clin Oncol*. 2007;25:1160–1168.
5. Conroy T, Galais M-P, Raoul JL, et al. UNICANCER-GI/FFCD PRODIGE Intergroup. Phase III randomized trial of definitive chemoradiotherapy (CRT) with FOLFOX or cisplatin and fluorouracil in esophageal cancer (EC): Final results of the PRODIGE 5/ACCORD 17 trial [abstract]. *J Clin Oncol*. 2012;30 (Suppl 18): LBA4003.
6. Khushalani NI, Leichman CG, Proulx G, et al. Oxaliplatin in combination with protracted-infusion fluorouracil and radiation: report of a clinical trial for patients with esophageal cancer. *J Clin Oncol*. 2002;20:2844–2850.
7. Lee SS, Kim SB, Park SI, et al. Capecitabine and cisplatin chemotherapy (XP) alone or sequentially combined chemoradiotherapy containing XP regimen in patients with three different settings of stage IV esophageal cancer. *Jpn J Clin Oncol*. 2007;37:829–835.
8. Javle MM, Yang G, Nwogu CE, et al. Capecitabine, oxaliplatin and radiotherapy: a phase 1B neoadjuvant study for esophageal cancer with gene expression analysis. *Cancer Invest*. 2009;27:193–200.
9. Sharma R, Yang GY, Nwogu CE et al. A single institution experience with neoadjuvant chemoradiation (CRT) with irinotecan (I) and cisplatin (C) in locally advanced esophageal carcinoma (LAEC) [abstract]. *J Clin Oncol*. 2009;27(Suppl 15): Abstract e15619.
10. Ajani JA, Winter K, Okawara GS, et al. Phase II trial of preoperative chemoradiation in patients with localized gastric adenocarcinoma (RTOG 9904): quality of combined modality therapy and pathologic response. *J Clin Oncol*. 2006;24:3953–3958.

*continued*

## References (continued)

11. Cunningham D, Allum WH, Stenning SP, et al. MAGIC Trial Participants. Perioperative chemotherapy versus surgery alone for resectable gastroesophageal cancer. *N Engl J Med*. 2006;355:11-20.
12. Sumpter K, Harper-Wynne C, et al. Report of two protocol planned interim analyses in a randomised multicentre phase III study comparing capecitabine with fluorouracil and oxaliplatin with cisplatin in patients with advanced oesophagogastric cancer receiving ECF. *Br J Cancer*. 2005;92:1976-1983.
13. Ychou M, Boige V, Pignon JP, et al. Perioperative chemotherapy compared with surgery alone for resectable gastroesophageal adenocarcinoma: an FNCLCC and FFCD multicenter phase III trial. *J Clin Oncol*. 2011;29:1715-1721.
14. Macdonald JS, Smalley SR, Benedetti J, et al. Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. *N Engl J Med*. 2001;345:725-730.
15. Jansen EP, Boot H, Saunders MP, et al. A phase I-II study of postoperative capecitabine-based chemoradiotherapy in gastric cancer. *Int J Radiat Oncol Biol Phys*. 2007;69:1424-1428.
16. André T, Quinaux E, Louvet C, et al. Phase III study comparing a semimonthly with a monthly regimen of fluorouracil and leucovorin as adjuvant treatment for stage II and III colon cancer patients: final results of GERCOR C96.1. *J Clin Oncol*. 2007;25:3732-3738.
17. Leong T, Joon DL, Willis D, et al. Adjuvant chemoradiation for gastric cancer using epirubicin, cisplatin, and 5-fluorouracil before and after three-dimensional conformal radiotherapy with concurrent infusional 5-fluorouracil: a multicenter study of the Trans-Tasman Radiation Oncology Group. *Int J Radiat Oncol Biol Phys*. 2011;79:690-695.
18. Lee HS, Choi Y, Hur WJ, et al. Pilot study of postoperative adjuvant chemoradiation for advanced gastric cancer: adjuvant 5-FU/cisplatin and chemoradiation with capecitabine. *World J Gastroenterol*. 2006;12:603-607.
19. Bang YJ, Kim YW, Yang HK, et al. CLASSIC trial investigators. Adjuvant capecitabine and oxaliplatin for gastric cancer after D2 gastrectomy (CLASSIC): a phase 3 open-label, randomised controlled trial. *Lancet*. 2012;379:315-321.
20. Lee J, Lim do H, Kim S, et al. Phase III trial comparing capecitabine plus cisplatin versus capecitabine plus cisplatin with concurrent capecitabine radiotherapy in completely resected gastric cancer with D2 lymph node dissection: the ARTIST trial. *J Clin Oncol*. 2012;30:268-273.
21. Bang YJ, Van Cutsem E, Feyereislova A, et al. ToGA Trial Investigators. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet*. 2010;376(9742):687-697.
22. Van Cutsem E, Moiseyenko VM, Tjulandin S, et al. V325 Study Group. Phase III study of docetaxel and cisplatin plus fluorouracil compared with cisplatin and fluorouracil as first-line therapy for advanced gastric cancer: a report of the V325 Study Group. *J Clin Oncol*. 2006;24:4991-4997.
23. Shah MA, Shibata S, Stoller RG, et al. MSKCC Gastric Cancer Consortium. Random assignment multicenter phase II study of modified docetaxel, cisplatin, fluorouracil (mDCF) versus DCF with growth factor support (GCSF) in metastatic gastroesophageal adenocarcinoma (GE). *J Clin Oncol*. 2010;28 (Suppl 15): 4010.
24. Shankaran V, Mulcahy MF, Hochster HS, et al. Docetaxel, oxaliplatin, and 5-fluorouracil for the treatment of metastatic or unresectable gastric or gastroesophageal junction (GEJ) adenocarcinomas: Preliminary results of a phase II study [abstract 47]. Presented at the 2009 Gastrointestinal Cancers Symposium.
25. Elkmern YM, Elsaid A, Al-Batran SE, et al. Final results of a phase II trial of docetaxel-carboplatin-FU in locally advanced gastric carcinoma. Presented at the Gastrointestinal Cancers Symposium, January 25-27, 2008, Orlando, FL.
26. Ross P, Nicolson M, Cunningham D, et al. Prospective randomized trial comparing mitomycin, cisplatin, and protracted venous-infusion fluorouracil (PVI 5-FU) with epirubicin, cisplatin, and PVI 5-FU in advanced esophagogastric cancer. *J Clin Oncol*. 2002;20:1996-2004.
27. Cunningham D, Starling N, Rao S, et al. Capecitabine and oxaliplatin for advanced esophagogastric cancer. *N Engl J Med*. 2008;358:36-46.
28. Lorenzen S, Schuster T, Porschen R, et al. Cetuximab plus cisplatin-5-fluorouracil versus cisplatin-5-fluorouracil alone in first-line metastatic squamous cell carcinoma of the esophagus: a randomized phase II study of the Arbeitsgemeinschaft Internistische Onkologie. *Ann Oncol*. 2009;20:1667-1673.
29. Al-Batran SE, Hartmann JT, Probst S, et al. Phase III trial in metastatic gastroesophageal adenocarcinoma with fluorouracil, leucovorin plus either oxaliplatin or cisplatin: a study of the Arbeitsgemeinschaft Internistische Onkologie. *J Clin Oncol*. 2008;26:1435-1442.
30. Kang YK, Kang WK, Shin DB, et al. Capecitabine/cisplatin versus 5-fluorouracil/cisplatin as first-line therapy in patients with advanced gastric cancer: a randomised phase III non-inferiority trial. *Ann Oncol*. 2009;20:666-673.
31. Bouché O, Raoul JL, Bonnetain F, et al. Fédération Francophone de Cancérologie Digestive Group. Randomized multicenter phase II trial of a biweekly regimen of fluorouracil and leucovorin (LV5FU2), LV5FU2 plus cisplatin, or LV5FU2 plus irinotecan in patients with previously untreated metastatic gastric cancer: a Federation Francophone de Cancérologie Digestive Group Study—FFCD 9803. *J Clin Oncol*. 2004;22:4319-4328.
32. Enzinger PC, Burntress B, Hollis D, et al. CALGB 80403/ECOG 1206: A randomized phase II study of three standard chemotherapy regimens (ECF, IC, FOLFOX) plus cetuximab in metastatic esophageal and GE junction cancer [abstract 4006]. *J Clin Oncol*. 2010; 28 (suppl 15):4007.
33. Kim GM, Jeung HC, Rha SY, et al. A randomized phase II trial of S-1-oxaliplatin versus capecitabine-oxaliplatin in advanced gastric cancer. *Eur J Cancer*. 2012;48:518-526.
34. Dank M, Zaluski J, Barone C, et al. Randomized phase III study comparing irinotecan combined with 5-fluorouracil and folinic acid to cisplatin combined with 5-fluorouracil in chemotherapy naive patients with advanced adenocarcinoma of the stomach or esophagogastric junction. *Ann Oncol*. 2008;19: 1450-1457.
35. Sym SJ, Hong J, Park J, et al. A randomized phase II study of biweekly irinotecan plus 5-fluorouracil/leucovorin (mFOLFIRI) in patients with metastatic gastric adenocarcinoma refractory to or progressive after first-line chemotherapy. *Cancer Chemother Pharmacol*. 2013;71:481-488.
36. Wolff K, Wein A, Reulbach U, et al. Weekly high-dose 5-fluorouracil as a 24-h infusion and sodium folinic acid (AIO regimen) plus irinotecan in patients with locally advanced nonresectable and metastatic adenocarcinoma or squamous cell carcinoma of the oesophagus: a phase II trial. *Anticancer Drugs*. 2009;20:165-173.
37. Ilson DH, Forastiere A, Arquette M, et al. A phase II trial of paclitaxel and cisplatin in patients with advanced carcinoma of the esophagus. *Cancer J*. 2000;6:316-323.
38. Petrasch S, Welt A, Reinacher A, et al. Chemotherapy with cisplatin and paclitaxel in patients with locally advanced, recurrent or metastatic oesophageal cancer. *Br J Cancer*. 1998;78:511-514.
39. Gadgeel SM, Shields AF, Heilbrun LK, et al. Phase II study of paclitaxel and carboplatin in patients with advanced gastric cancer. *Am J Clin Oncol*. 2003;26:37-41.

## References (continued)

40. Ajani JA, Fodor MB, Tjulandin SA, et al. Phase II multi-institutional randomized trial of docetaxel plus cisplatin with or without fluorouracil in patients with untreated, advanced gastric, or gastroesophageal adenocarcinoma. *J Clin Oncol*. 2005;23:5660-5667.
41. Kim JY, Do YR, Park KU, et al. A multi-center phase II study of docetaxel plus cisplatin as first-line therapy in patients with metastatic squamous cell esophageal cancer. *Cancer Chemother Pharmacol*. 2010;66:31-36.
42. Burtness B, Gibson M, Egleston B, et al. Phase II trial of docetaxel-irinotecan combination in advanced esophageal cancer. *Ann Oncol*. 2009;20:1242-1248.
43. Ohtsu A, Shimada Y, Shirao K, et al. Randomized phase III trial of fluorouracil alone versus fluorouracil plus cisplatin versus uracil and tegafur plus mitomycin in patients with unresectable, advanced gastric cancer: The Japan Clinical Oncology Group Study (JCOG9205). *J Clin Oncol*. 2003;21:54-59.
44. Hong YS, Song SY, Lee SI, et al. A phase II trial of capecitabine in previously untreated patients with advanced and/or metastatic gastric cancer. *Ann Oncol*. 2004;15:1344-1347.
45. Albertsson M, Johansson B, Friesland S, et al. Phase II studies on docetaxel alone every third week, or weekly in combination with gemcitabine in patients with primarily locally advanced, metastatic, or recurrent esophageal cancer. *Med Oncol*. 2007;24:407-412.
46. Ford HE, Marshall A, Bridgewater JA, et al. Docetaxel versus active symptom control for refractory oesophago-gastric adenocarcinoma (COUGAR-02): an open-label, phase 3 randomised controlled trial. *Lancet Oncol*. 2014;15:78-86.
47. Ajani JA, Ilson DH, Daugherty K, et al. Activity of taxol in patients with squamous cell carcinoma and adenocarcinoma of the esophagus. *J Natl Cancer Inst*. 1994;86:1086-1091.
48. Ilson DH, Wadleigh RG, Leichman LP, Kelsen DP. Paclitaxel given by a weekly 1-h infusion in advanced esophageal cancer. *Ann Oncol*. 2007;18:898-902.
49. Fuchs CS, Tomasek J, Yong CJ, et al. Ramucirumab monotherapy for previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (REGARD): an international, randomized, multicentre, placebo-controlled, phase 3 trial. *Lancet*. 2014;383:31-39.
50. Wilke H, Van Cutsem E, Oh SC, et al. RAINBOW: A global, phase III, randomized, double-blind study of ramucirumab plus paclitaxel versus placebo plus paclitaxel in the treatment of metastatic gastroesophageal junction (GEJ) and gastric adenocarcinoma following disease progression on first-line platinum- and fluoropyrimidine-containing combination therapy rainbow IMCL COP12-0922 (I4T-IE-JVBE). *J Clin Oncol*. 2014;32 (3\_suppl):Abstract LBA7.
51. Hironaka S, Ueda S, Yasui H, et al. Randomized, Open-Label, Phase III Study Comparing Irinotecan With Paclitaxel in Patients With Advanced Gastric Cancer Without Severe Peritoneal Metastasis After Failure of Prior Combination Using Fluoropyrimidine Plus Platinum: WJOG 4007 Trial. *J Clin Oncol*. 2013;31:4438-4444.
52. Thuss-Patience PC, Kretzschmar A, Bichev D, et al. Survival advantage for irinotecan versus best supportive care as second-line chemotherapy in gastric cancer—a randomized phase III study of the Arbeitsgemeinschaft Internistische Onkologie (AIO). *Eur J Cancer*. 2011;47:2306-2314.
53. Fuchs CS, Moore MR, Harker G, et al. Phase III comparison of two irinotecan dosing regimens in second-line therapy of metastatic colorectal cancer. *J Clin Oncol*. 2003;21:807-814.
54. Ilson DH. Phase II trial of weekly irinotecan/cisplatin in advanced esophageal cancer. *Oncology (Williston Park)*. 2004;18:22-25.
55. Leary A, Assersohn L, Cunningham D, et al. A phase II trial evaluating capecitabine and irinotecan as second line treatment in patients with oesophago-gastric cancer who have progressed on, or within 3 months of platinum-based chemotherapy. *Cancer Chemother Pharmacol*. 2009;64:455-462.
56. Maugeri-Sacca M, Pizzuti L, Sergi D, et al. FOLFIRI as a second-line therapy in patients with docetaxel-pretreated gastric cancer: a historic cohort. *J Exp Clin Cancer Res*. 2013;32:67.
57. Lustberg MB, Bekali-Saab T, Young D, et al. Phase II randomized study of two regimens of sequentially administered mitomycin C and irinotecan in patients with unresectable esophageal and gastroesophageal adenocarcinoma. *J Thorac Oncol*. 2010;5:713-718.
58. Giuliani F, Molica S, Maiello E, et al. Irinotecan (CPT-11) and mitomycin-C (MMC) as second-line therapy in advanced gastric cancer: a phase II study of the Gruppo Oncologico dell' Italia Meridionale (prot. 2106). *Am J Clin Oncol*. 2005;28:581-585.
59. Barnias A, Papamichael D, Syrigos K, Pavlidis N. Phase II study of irinotecan and mitomycin C in 5-fluorouracil-pretreated patients with advanced colorectal and gastric cancer. *J Chemother*. 2003;15:275-281.
60. Hofheinz RD, Hartung G, Samel S, et al. High-dose 5-fluorouracil/folinic acid in combination with three-weekly mitomycin C in the treatment of advanced gastric cancer: A phase II study. *Onkologie*. 2002;25:255-260.